



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY
WASHINGTON, D.C. 20460

WELL FILE
Casefile
3232

OFFICE OF
PESTICIDES AND TOXIC SUBSTANCES

MEMORANDUM

SUBJECT: Dietary Exposure Analysis for the Proposed Use of Vinclozolin (Ronilan) on Greenhouse-grown Tomatoes Imported from Spain and the Netherlands, 8E3688

FROM: J. Robert Tomerlin, Ph.D. *J.R. Tomerlin* 4/28/89
Tolerance Assessment System Staff
HED/SACB (H7509C)

THROUGH: R. Bruce Jaeger *RBJ 4/28/89*
Head, Special Analysis and Outreach Section
HED/SACB (H7509C)

TO: Susan Lewis, PM 21
Herbicide-Fungicide Branch
Registration Division (H7505C)

Action Requested

Provide an estimate of dietary exposure resulting from the proposed use of vinclozolin on imported greenhouse-grown cucumbers and tomatoes.

Discussion

1. Toxicology Endpoint: The routine chronic TAS analysis used a reference dose (ADI) of 0.025 mg/kg body weight/day, based upon a NOEL of 2.5 mg/kg body weight/day and an uncertainty factor of 100 from a 6 month dog feeding study. This value has been approved by HED (5/19/86) and Agency (7/8/86) reference dose committees.

2. Residue Information: Food uses evaluated were published tolerances from 40 CFR 180.380 and the proposed tolerances on cucumbers and tomatoes (W. T. Chin memo, 4/27/89). A summary of the residue information used in the analysis is attached as Table 1.

3. Exposure Analysis: The TAS chronic exposure analysis uses tolerance level residues and 100 per cent crop treated to estimate the Theoretical Maximum Residue Contribution (TMRC) for the overall U.S. population and 22 population subgroups. The estimated TMRC for the overall U.S. population was calculated as 0.018815 mg/kg body weight/day, which represents 75% of the ADI. The two most highly exposed TAS subgroups, nursing and non-nursing infants, had TMRCs of 0.044215 mg/kg body weight/day (177% of the ADI) and

Vinclozolin Dietary Exposure Analysis, page 2

0.089879 mg/kg body weight/day (360% of the ADI), respectively. A complete TMRC summary is shown in Table 2.

Since there are other pending actions in the data file, the "DIFFERENCE..." column in Table 2 gives the increase in exposure, in terms of the per cent of the ADI, resulting from the proposed tolerances on cucumbers and tomatoes.

4. Comments: The bulk of the exposure for the three TAS subgroups discussed comes from published tolerances. The actual exposure is likely lower than what is estimated in this analysis because we used tolerance level residues and assume that 100 per cent of the crops were treated. An exposure estimate closer to actual exposure could be calculated if anticipated residue and per cent crop treated data were available for stone fruits, particularly peaches. Peaches account for over 20% of the ADI for the overall U.S. population, over 130% of the ADI for nursing infants, and over 200% of the ADI for non-nursing infants.

The exposure resulting from the proposed tolerances on imported greenhouse-grown cucumbers and tomatoes is also overestimated, because all tomatoes consumed were assumed to have been treated with vinclozolin, regardless of country of origin or type of production.

Attachments

cc: TAS (Tomerlin, SACB), DEB, Caswell #323C, B. Jaeger,
Anderson (TOX-HAFS)

Table 1

CHEMICAL INFORMATION FOR CASWELL NUMBER 323C

DATE: 04/27/89 PAGE: 1

CHEMICAL	STUDY TYPE	EFFECTS	REFERENCE DOSES	DATA GAPS/COMMENTS	STATUS
Vinclozolin (Ronilan) Caswell #123C CAS No. 50471-44-8 A.I. CODE: 113201 CFR No. 180.380	6mo feeding- dog NOEL= 2,500.0 mg/kg LEL= 100.00 ppm LEL= 7,500.0 mg/kg ONCO: Class E (HED WOTE).	Increased absolute and relative adrenal weights. Histo changes seen in adrenal at higher doses. No evidence of oncogenicity in rats or mice.	ADI SF -->100 OPP RfD= 0.025000 EPA RfD= 0.025000	no data gaps. WHO last reviewed 1988; temp. ADI established. On IRIS.	HED complete 05/19/86. EPA verified 07/08/86.

FOOD CODE	FOOD NAME	PETITION NUMBER	NEW	TOLERANCE (PPM)	PUBLISHED
01006AA	RASPBERRIES	3F2934		10.0000	
01014AA	GRAPE-FRESH	1E2457		6.0000	
01014DA	GRAPE-RAISINS	7H5529		30.0000 H	
01014JA	GRAPE-JUICE	1E2457		6.0000	
01016AA	STRAWBERRIES	9F7205		10.0000	
05001AA	APRICOTS-FRESH	2F2650		25.0000	
05001DA	APRICOTS-DRIED	2F2650		25.0000	
05002AA	CHEERIES-FRESH	2F2650		25.0000	
05002DA	CHEERIES-DRIED	2F2650		25.0000	
05002JA	CHEERIES-JUICE	2F2650		25.0000	
05003AA	NECTARINES	2F2650		25.0000	
05004AA	PEACHES-FRESH	2F2650		25.0000	
05004DA	PEACHES-DRIED	2F2650		25.0000	
05005AA	PLUMS(DAMSONS)-FRESH	2F2650		25.0000	
05005DA	PLUMS-PRUNES(DRIED)	5H5462		75.0000 H	
05005JA	PLUMS/PRUNE-JUICE	5H5462		75.0000 H	
06018RA	KIWI	0E2380		10.0000	
10010RA	CUCUMBERS	8E3688	1.0000		
11001RA	PEPPERS-SWEET, GARDEN	4E2998		3.0000	
11002AA	TOMATOES-WHOLE	8E3688		3.0000	
11005AA	TOMATOES-JUICE	8E3688		3.0000	
11005RA	TOMATOES-PUREE	8E3688		3.0000	
11005TA	TOMATOES-PASTE	8E3688		3.0000	
11005UA	TOMATOES-CATSUP	8E3688		3.0000	
13003RA	CHICORY (FRENCH OR BELGIAN ENDIVE)	8E3620		5.0000	
13013AA	LETUCE-LEAFY VARIETIES	3F2934		10.0000	
13020RA	LETUCE-UNSPECIFIED	2F2595		10.0000	
13045AA	LETUCE-HEAD VARIETIES	2F2595		10.0000	
14007AA	GARLIC	3F2934		1.0000	
14011AA	ONIONS-DRY-BULB (CIPOLLINI)	3F2934		1.0000	
14011DA	ONIONS-DEHYDRATED OR DRIED	3F2934		1.0000	
43058AA	WINE AND SHERRY	1E2457		6.0000	

J

Table 2

TOLERANCE ASSESSMENT SYSTEM ROUTINE CHRONIC ANALYSIS

DATE: 04/27/89

PAGE: 1

CHEMICAL INFORMATION	STUDY TYPE	EFFECTS		REFERENCE DOSES	DATA GAPS/COMMENTS	STATUS	
		Increased absolute and relative adrenal weights.	Histo changes seen in adrenal at higher doses.				
Vinclozolin (Ronilan)	6mo feeding - dog NOEL= 2.5000 mg/kg 100.00 ppm			ADI SF -->100 OPP RFD= 0.025000 EPA RFD= 0.025000	No data gaps.	HED complete 05/19/86. EPA verified 07/08/86.	
Caswell #323C	LEL= 7.5000 mg/kg 300.00 ppm	Histo changes seen in adrenal at higher doses.				WHO last reviewed 1988; temp. ADI established.	
CAS No. 50471-44-8		No evidence of oncogenicity in rats or mice.				On IRIS.	
A.I. CODE: 113201							
CFR No. 180.380	ONCO: Class E (HED NOTE).						
TOTAL TMRC (MG/KG BODY WEIGHT/DAY)		NEW TMRC**		NEW TMRC AS PERCENT OF RFD		EFFECT OF ANTICIPATED RESIDUES	
POPULATION SUBGROUP	CURRENT TMRC*	CURRENT TMRC*	NEW TMRC**	DIFFERENCE AS PERCENT OF RFD	ARC	ARC	%RFD
U.S. POPULATION - 48 STATES	0.014378	0.018815	75.258160	17.745160			
U.S. POPULATION - SPRING SEASON	0.014570	0.018774	75.095844	16.815804			
U.S. POPULATION - SUMMER SEASON	0.019597	0.024233	96.933640	18.545160			
U.S. POPULATION - FALL SEASON	0.011811	0.016255	65.021036	17.775296			
U.S. POPULATION - WINTER SEASON	0.011549	0.016012	64.048696	17.851196			
NORTHEAST REGION	0.015894	0.020461	81.844336	18.269612			
NORTH CENTRAL REGION	0.014456	0.019121	76.482044	18.659552			
SOUTHERN REGION	0.010556	0.014529	58.115240	15.890680			
WESTERN REGION	0.018858	0.023588	94.350000	18.917620			
HISPANICS	0.016187	0.021376	85.505376	20.758492			
NON-HISPANIC WHITES	0.015065	0.019671	78.585984	18.424440			
NON-HISPANIC BLACKS	0.009284	0.012333	49.333812	12.198116			
NON-HISPANIC OTHERS	0.012645	0.016505	66.019136	15.440612			
NURSING INFANTS (< 1 YEAR OLD)	0.043414	0.044215	176.860764	3.203156			
NON-NURSING INFANTS (< 1 YEAR OLD)	0.086940	0.098879	359.516964	11.757980			
FEMALES (13+ YEARS, PREGNANT)	0.012391	0.016333	65.330463	15.767148			
FEMALES 13+ YEARS, NURSING	0.014154	0.018208	72.833748	16.219728			
CHILDREN (1-6 YEARS OLD)	0.028640	0.037166	148.663276	34.104012			
CHILDREN (7-12 YEARS OLD)	0.017718	0.024599	98.397964	27.524328			
MALES (13-19 YEARS OLD)	0.008719	0.013617	54.467636	19.593352			
FEMALES (13-19 YEARS OLD, NOT PREG. OR NURS.)	0.009298	0.013381	53.522968	16.330116			
MALES (20 YEARS AND OLDER)	0.010252	0.013796	55.183332	14.174668			
FEMALES (20 YEARS AND OLDER, NOT PREG. OR NURS.)	0.012353	0.015752	63.009352	13.597012			

*Current TMRC does not include new or pending tolerances.
**New TMRC includes new, pending, and published tolerances.

4